

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Flicca (Stone, Bird Migration Records of William Bartram, Auk, xxx, 1913, 341).

Flying Auger, Martha's Vineyard, Mass. (McAtee, Local Names of Water Fowl and Other Birds, p. 19).

Golden-Headed Woodpecker or Yellowhammer (Hurst's New Nuttall's Dictionary, Suppli., p. 871). Citation by Miller, Oologist, xxxi, p. 10. A book name that I was inclined to reject as a typographical error, but after all, it is no more ridiculous than some other names in current use.

Graywacker, Eastern Shore of Maryland (Oberholser), (McAtee, *Ibid.*).

Looping Bird, Church's Island, N. C. (McAtee, Ibid.).

Rampike, Newfoundland (Montreal Family Herald and Weekly Star, Oct. 18, 1911). For which I am under obligations to Mr. McAtee. F. L. Burns.

Berwyn, Pa.

THE NEST LIFE OF THE WESTERN HOUSE WREN. (Abstract.)

Read before Wilson Ornithological Club, third annual meeting, Columbus, Ohio, 1915.

BY ALTHEA R. SHERMAN.

Studies of the behavior outside of the nest of *Troglodytes aëdon* parkmani were made for the most part from birds nesting in boxes in various places in the yard, and the inner activities of its home were studied through peepholes that commanded views of the inside of nests, situated in a barn. From these vantage points was watched the progress of the nests from their building, until they were deserted by the young. The duration of the nest period, as is the case with other species, is subject to considerable variation, the average time for nest occupation, counting from the depositing of the first egg, has been found to be thirty-two days.

There has existed among ornithologists a general misconception relating to the amount of light the nest receives through the hole. A mere glance into the boxes provided with peep-holes is sufficient to convince any one that the illumination of the nest is ample, and differs little in intensity from the lighting of most of our living-rooms.

To feed and care for all the nestlings alike is the normal conduct of a bird, and the abnormal happens, when a portion of a brood thrives vigorously, while others die from starvation. Deaths from starving in the nest of the Western House Wren were noted to result from causes very different from those that brought death to the young of the Northern Flicker.